

小鼠胚胎干细胞 J1 说明书

目录号: SCSP-219

细胞名称: J1

细胞描述: J1 细胞来源于雄性鼠灰色的 129S4/SvJae 胚胎。该细胞能进入生殖系。

物种: 小鼠, 129S4/SvJae 品系, 雄性

细胞来源: 2013 年引进

组织: 内细胞团

生物安全等级: BSL-1

完全培养液配方: mES 完全培养液 (产品编号 SCSP-603)。配方见备注。

冻存日期/代数: 详见 冻存管/培养瓶 标识

参考传代周期: 2–3 天

参考传代比例: 1:4–1:7

参考换液频率: 每天

冻存液配方: 培养液 80%, FBS 10%, DMSO 10%

细胞状态: 球形克隆

支原体检测结果: 阴性

参考文献:

Stevens LC. A new inbred subline of mice (129-terSv) with a high incidence of spontaneous congenital testicular teratomas. *J. Natl. Cancer Inst.* 50: 235–242, 1973. PubMed: 4692863

Li E, et al. Targeted mutation of the DNA methyltransferase gene results in embryonic lethality. *Cell* 69: 915–926, 1992. PubMed: 1606615

Krzyzanowski PM, et al. Identification of novel stem cell markers using gap analysis of gene expression data. *Genome Biology* 8(9): R193, 2007. PubMed: 17875203

Luikenhuis S, et. al. Antisense transcription through the Xist locus mediates Tsix function in embryonic stem cells. *Mol. Cell Biol.* 21(24): 8512–8520, 2001. PubMed: 11713286

Goto T, et al. Regulation of X-Chromosome inactivation in development in mice and humans. *Micro. Molec. Biol. Rev. (ASM)* 62(2): 362–378, 1998. PubMed: 9618446

Yap DYL, et al. Using biomarker signature patterns for an mRNA molecular diagnostic of mouse embryonic stem cell differentiation state. *BMC Genomics* 8: 210, 2007. PubMed: 17605829

Haase VH, et al. Vascular tumors in livers with targeted inactivation of the von Hippel-Lindau tumor suppressor. *Proc. Natl. Acad. Sci.* 98(4): 1583–1588, 2001. PubMed: 11171994

Oda M, et al. DNA methylation regulates long-range gene silencing of an X-linked homeobox gene cluster in a lineage-specific manner. *Genes Dev.* 20(24): 3382?3394, 2006.
PubMed: 17182866

Lorincz MC, et al. DNA methylation density influences the stability of an epigenetic imprint and Dnmt3a/b-independent de novo methylation. *Mol. Cell Biol.* 22(21): 7572?7580, 2002. PubMed: 12370304

Tucker KL, et al. A transgenic mouse strain expressing four drug-selectable marker genes. *Nucleic Acids Res.* 25: 3745-3746, 1997. PubMed: 9278500

Biniszkiewicz D, et al. Dnmt1 overexpression causes genomic hypermethylation, loss of imprinting, and embryonic lethality. *Mol. Cell Biol.* 22(7): 2124?2135, 2002. PubMed: 11884600

备注:

1. mES 完全培养液配方 (600 ml) :

DMEM (Invitrogen, 111960-044)	497 ml
ES 级 FBS (Gibco)	90 ml
Glutamax (Invitrogen, 35050)	6 ml
NEAA (Invitrogen, 11140)	6 ml
LIF (Millipore, ESG1107)	60 μ l (终浓度为 1000U/ml)
β -Mer (Invitrogen, 21985)	1.5 ml

2. 相关产品: J1 所使用的饲养层细胞: CF-1 MEF

本细胞库可提供:

SCSP-105 P0,可以增殖

SCSP-105R 已经经过射线处理, 不再增殖, 可直接作为 feeder 使用。

3. 我库冻存时, 体积为 500 μ l, 预期存活率 70%, 每支冻存管的细胞复苏, 3 至 4 天后, 会形成 30 至 40 个克隆, 建议复苏至 1 个 T25 培养瓶中。

中国科学院典型培养物保藏委员会细胞库/干细胞库